

Balanced CoQ₆ biosynthesis is required for lifespan and mitophagy in yeast

I. González-Mariscal, A. Martín-Montalvo, C. Ojeda-González, A. Rodríguez-Eguren, P. Gutiérrez-Ríos, P. Navas, and C. Santos-Ocaña*

Centro Andaluz de Biología del Desarrollo, Universidad Pablo de Olavide-CSIC, CIBERER Instituto de Salud Carlos III, Sevilla, 41013, Spain.

* Corresponding Author:

C. Santos-Ocaña, E-mail: csanoca@upo.es

ABSTRACT Coenzyme Q is an essential lipid with redox capacity that is present in all organisms. In yeast its biosynthesis depends on a multiprotein complex in which Coq7 protein has both catalytic and regulatory functions. Coq7 modulates CoQ₆ levels through a phosphorylation cycle, where dephosphorylation of three amino acids (Ser/Thr) by the mitochondrial phosphatase Ptc7 increases the levels of CoQ₆. Here we analyzed the role of Ptc7 and the phosphorylation state of Coq7 in yeast mitochondrial function. The conversion of the three Ser/Thr to alanine led to a permanently active form of Coq7 that caused a 2.5-fold increase of CoQ₆ levels, albeit decreased mitochondrial respiratory chain activity and oxidative stress resistance capacity. This resulted in an increase in endogenous ROS production and shortened the chronological life span (CLS) compared to wild type. The null *PTC7* mutant (*ptc7Δ*) strain showed a lower biosynthesis rate of CoQ₆ and a significant shortening of the CLS. The reduced CLS observed in *ptc7Δ* was restored by the overexpression of *PTC7* but not by the addition of exogenous CoQ₆. Overexpression of *PTC7* increased mitophagy in a wild type strain. This finding suggests an additional Ptc7 function beyond the regulation of CoQ biosynthesis. Genetic disruption of *PTC7* prevented mitophagy activation in conditions of nitrogen deprivation. In brief, we show that, in yeast, Ptc7 modulates the adaptation to respiratory metabolism by dephosphorylating Coq7 to supply newly synthesized CoQ₆, and by activating mitophagy to remove defective mitochondria at stationary phase, guaranteeing a proper CLS in yeast.

doi: 10.15698/mic2017.02.556

Received originally: 19.07.2016;

in revised form: 29.11.2016,

Accepted 19.01.2017,

Published 03.02.2017.

Keywords: coenzyme Q₆, regulation, mitochondria, yeast, mitophagy, chronological life span.

Abbreviations:

CLS - chronological life span.

INTRODUCTION

Coenzyme Q (CoQ) deficiency is a syndrome that belongs to the family of mitochondrial diseases [1]. CoQ, a lipid embedded in cell membranes, main function is to act as an electron carrier and as an antioxidant. CoQ deficiency is classified by the observed phenotype as it is a multiple-caused syndrome [2]. There are primary and secondary CoQ deficiencies; the deficiency can be a consequence of mutations in genes involved directly in the CoQ biosynthesis (primary deficiency) [3, 4] or a consequence of defects or mutations in genes not directly related to CoQ biosynthesis (secondary deficiency) [5, 6]. The existence of secondary deficiencies introduces the idea of regulatory mechanisms of CoQ biosynthesis that could also be related to a general regulation of mitochondrial metabolism. The main defect caused by CoQ deficiency is a depletion of ATP

in tissues. The variety of symptoms and the diversity of CoQ functions introduce a source of complexity in the analysis of CoQ deficiency syndrome that requires the analysis of the mechanisms regulating the biosynthesis of CoQ [7].

In yeast, CoQ₆ is synthesized in the mitochondria after the condensation of an activated polyisoprenoid tail with a hydroxybenzoic ring; the ring is further modified by several reactions [8]. The enzymes that catalyze these reactions are encoded by nuclear *COQ* genes [9]. Most of the *COQ* genes encode proteins responsible for enzymatic activity; however, three proteins without enzymatic activity, Coq4, Coq8 and Coq9, play a structural role. Coq4 has been reported to support the assembly of the CoQ₆ biosynthetic complex in yeast [10]. Coq8 has been included in a family




FIGURE 1: CoQ₆-dependent mitochondrial respiratory chain activities are negatively affected in yeast expressing Coq7 phosphomutant versions. Mitochondrial activities were measured with mitochondria purified from the *coq7Δ* strain harboring the indicated plasmids. pNMQ7 corresponds to the wild type *COQ7* gene, pRS316 corresponds to the empty vector, pAAA and pDED correspond to the Coq7 phosphomutant versions and pmQ7 corresponds to a multicopy expression of wild type *COQ7* gene. **(A)** Top panel: CoQ₆ quantification, bottom panel Western blots of mitochondrial samples probed with antibodies against Coq7. Mitochondrial activities: **(B)** NADH-DCIP reductase, **(C)** Complex II: succinate-DCIP reductase, **(D)** Complex III: Decylubiquinol-cytochrome c reductase, **(E)** NADH-cytochrome c reductase and **(F)** Complex II+III: Succinate-cytochrome c reductase. Results are expressed as nmol/mg mitochondrial protein·min. Data are mean ± SD, N ≥ 3 independent assays. ** P ≤ 0.001 compared to positive control samples.

of unusual kinases [11], which also includes other proteins involved in CoQ biosynthesis [12]. Recent analysis of the function of human ADCK3 protein, homologous of yeast Coq8, showed that the inhibition of ADCK3 kinase activity is required for the activation of CoQ₆ biosynthesis [13, 14]. Coq9 is a membrane protein located at the matrix side of the mitochondrial inner membrane and it belongs to the CoQ₆ biosynthetic complex, where it co-migrates with Coq3 and Coq4 at a molecular mass of approximately 1 MDa [15] and it binds to Coq7 to promote CoQ₆ biosynthesis [16].

Yeast biosynthesis of CoQ₆ occurs in a multi-protein complex (Q-synthome). The assembly of the Q-synthome requires the post-translational modification of Coq proteins. Several studies in the last years have demonstrated the existence of the Q-synthome [17–19] and several models for the assembly of the complex have been proposed [9, 20]. The complex assembly starts with a nucleation around the quinone-like lipid polyisoprenyl benzoate bound to a nucleating Coq protein such as Coq4. The nucleation step is ended with the assembly of a pre-complex that accumulates a CoQ₆ intermediate, the demethoxy quinone (DMQ₆) [19, 21]. DMQ₆ is converted to CoQ₆ after the activation of Coq7 by dephosphorylation [22]. Coq7 catalyzes the next to last reaction of the pathway [23], the DMQ₆ hydroxylation. Several studies have reported the existence of phosphoproteins in the family of Coq proteins: Coq3, Coq5 and Coq7 [24, 25], but only phosphorylation of Coq7 is known to have a physiological relevance [22]. Coq7 phosphorylation leads to a low activity state, therefore accumulating DMQ₆, while its dephosphorylation activates Coq7 and increases CoQ₆ levels. Both activation states of Coq7 can be achieved by changing the carbon source in the culture media [22]. These results were confirmed in *COQ7* null mutants yeast strains (*coq7Δ*) expressing either a Coq7 version that is permanently dephosphorylated (Coq7-AAA), associated with a sharp increase of CoQ₆ concentration, or a permanently phosphorylated version (Coq7-DED) that is associated with a significant decrease of CoQ₆ levels [22]. The Coq7-AAA version was obtained by site-directed mutagenesis of residues S20, S28 and T32 to alanine, while in the Coq7-DED version these residues were mutated to glutamic or aspartic acid. Recent studies have demonstrated the presence of another phosphorylatable residue in Coq7, the S114 [26], whose modification affects the catalytic function of Coq7. The activation (i.e. dephosphorylation) of Coq7 is carried out by Ptc7, a phosphatase that belongs to the type 2C Mg²⁺ or Mn²⁺ dependent protein phosphatases, PPM [27, 28]. Other mitochondrial members of this family (Ptc5) have been related to the activation of pyruvate dehydrogenase (PDH) [29, 30] and with the activation of mitophagy (Ptc6) [31, 32]. Null mutants of *PTC7* gene (*ptc7Δ*) have decreased levels of CoQ₆, decreased mitochondrial respiratory chain activities and decreased resistance to oxidative stress [28]. The expression of Coq7-AAA in *ptc7Δ* did not disrupt the high amount of CoQ₆ produced by this Coq7 version, demonstrating a relationship between Coq7 and Ptc7 [28].

Here we investigate the effect of both Coq7 mutants, Coq7-AAA and Coq7-DED on mitochondrial physiology and

its relationship with Ptc7 function. Our results indicate that although Coq7 mutants modify the mitochondrial physiology in a similar fashion than *ptc7Δ* relative to CoQ₆ levels, they have different effects on chronological life span, respiratory complexes interactions and on mitophagy activation.

RESULTS

Coq7-phosphorylation mutants show defects in mitochondrial respiratory chain activities

We measured CoQ₆ content and the activities of mitochondrial respiratory chain (MRC) in single or coupled complexes (Figure 1). The strain *coq7Δ* did not contain CoQ₆, which was rescued in the control strain (*coq7Δ/pNMQ7*). The strain expressing permanently dephosphorylated Coq7 (*coq7Δ/pAAA*) showed a dramatic increase of CoQ₆, while the strain expressing permanently phosphorylated Coq7 (*coq7Δ/pDED*) shows a significant decrease of CoQ₆ compared to control. Multicopy *COQ7* transformed yeast (*coq7Δ/pmQ7*) also significantly increased CoQ₆ (Figure 1A). Under the experimental conditions and in all the strains analyzed, Coq7 was detected by western blotting at the expected size of 24 kDa. NADH-Q reductase activity, measured as NADH-DCIP reductase, was decreased in the *coq7Δ/pDED* strain in a similar manner than negative control (*coq7Δ/pRS316*), but in the *coq7Δ/pAAA* strain, the activity was increased over the control, equivalent to the activity measured in the *coq7Δ/pmQ7* strain (Figure 1B). Complex II activity measured as succinate-DCIP reductase showed a moderated decrease in both *coq7Δ/pDED* and *coq7Δ/pAAA* strains (Figure 1C). However, complex II activity was increased significantly in the *coq7Δ/pmQ7* strain. Complex III activity (decydubiquinol-cytochrome c reductase) showed changes comparable to those in complex I (Figure 1D). Coupled MRC activities require CoQ₆ as electron carrier, which is not added exogenously in the assay. Complexes activities such as NADH-cytochrome c reductase and succinate-cytochrome c reductase (Figures 1E and 1F) were decreased in both *coq7Δ/pAAA* and *coq7Δ/pDED* strains compared to control. Interestingly, the decrease in NADH-cytochrome c reductase activity in the *coq7Δ/pAAA* strain does not correlate with the increased activity observed in single complexes (Figure 1B and 1D) and with the high amount of CoQ₆ found in mitochondria of the *coq7Δ/pAAA* strain. The activities of these complexes in the *coq7Δ/pmQ7* strain were significantly higher than in control.

Oxidative stress conditions in Coq7 phosphomutants

Due to the changes observed in MRC, we analyzed the endogenous oxidative stress, measured as H₂O₂ generation in mitochondria, from *coq7Δ* yeast expressing the different versions of Coq7 (Figure 2A). Expression of both Coq7-pAAA and Coq7-pDED showed an increased generation of H₂O₂ in mitochondria compared to control, while the *coq7Δ/pmQ7* strain showed a decreased amount, even lower than control. To determine the oxidative stress pro-




FIGURE 2: Expression of Coq7 phosphomutant versions increases both the endogenous generation and the sensitivity to oxidative stress. (A) Quantification of hydrogen peroxide generation was performed in presence of NADH and succinate as electron donors and 50 µg of purified mitochondria from the indicated strains. Data are mean ± SD, N ≥ 3 independent assays. (B) Superoxide anion generation was measured using acetylated cytochrome c as electron acceptor. Quantification was produced in presence of reduced decylubiquinone as electron donor and 50 µg of purified mitochondria from the indicated strains. Data are mean ± SD, N ≥ 3 independent assays. ** P ≤ 0.001 compared to positive control (*coq7Δ/pNMQ7*). (C) Oxidative stress sensitivity was measured in the indicated strains. Cells were subjected to oxidative stress treatment by H_2O_2 (0.75 mM) or linolenic acid (1 mM) in 0.1 M sodium phosphate buffer plus 0.2 % glucose buffer at 150×10^6 cells/ml. At the indicated times cells were harvested, diluted in sterile water and seeded in triplicate in YPD plates to calculate the number of CFU. The experiment is representative of a set of three. (D) Cell viability was monitored under induced oxidative stress. Yeast cells were grown in SDc –ura glucose. After two days, cells were spotted by serial dilutions (1/10) from 0.5 OD_{660nm}/ml onto YPD plates containing the indicated concentrations of oxidative stress agents. Plates were incubated at 30°C for 3 days and imaged.

duced specifically by the complex III, we analyzed the superoxide anion generation using reduced decylubiquinone and acetylated cytochrome *c* (Figure 2B) [33]. Strains expressing both mutated versions of Coq7 produced significantly higher amounts of superoxide, from 200 to 400%, compared to wild type. Also, superoxide was higher in the *coq7Δ/pRS316* strain. On the contrary, the *coq7Δ/pmQ7* strain showed a superoxide production comparable to control.

In yeast, CoQ₆ acts as a powerful antioxidant and it is required to protect against lipophilic oxidants such as linolenic acid [34–36]. The sensitivity to either H₂O₂ or linolenic acid was analyzed in a *coq7Δ* strain expressing both Coq7 versions harboring phosphosite modifications (Figures 2C–D). Survival was not compromised in all strains without oxidative stress insult (Figure 2C) but after incubation with H₂O₂ or linolenic acid the wild type strain showed the higher survival rate at 2, 4 and 6 hours, while the negative control (*coq7Δ/pRS316*) showed a lower survival rate. Interestingly, both treatments in *coq7Δ/pAAA* and *coq7Δ/pDED* strains produced a similar effect on survival, being higher than negative control but lower than control. A similar effect was found when the assay was performed in agar plates with H₂O₂ (Figure 2D); the lack of CoQ₆ in the *coq7Δ/pRS316* strain compromised the growth but *coq7Δ/pAAA* and *coq7Δ/pDED* growth was only slightly lower compared to control. When the assay was performed with linolenic acid at higher concentration, both *coq7* mutants, *coq7Δ/pAAA* and *coq7Δ/pDED*, were more sensitive to the treatment, although both retain CoQ₆ production. That result is surprising since null Coq mutants do not produce CoQ₆ and show a high sensitivity to linolenic acid [35, 36]. In contrast, *coq7Δ/pAAA* and *coq7Δ/pDED* strains synthesize CoQ₆, with higher production than control for *coq7Δ/pAAA* strain.

Chronological life span is differentially affected in Coq7 phosphomutants

Chronological lifespan (CLS) measurement refers to yeast cells longevity in an exhausted culture media after the onset of stationary phase [37]. The extension of CLS depends on the existence of an intact respiratory metabolism [38–41] and also by the expression of intact antioxidant defenses [42–44]. Given that these requirements are affected in cells expressing both Coq7-pAAA and Coq7-pDED, CLS have been measured in these cells (Figure 3A and Table S1). The *coq7Δ/pRS316* strain showed shorter mean CLS (2.8 ± 0.2 days) compared to both *coq7Δ/pNMQ7* (12.2 ± 0.7 days) and *coq7Δ/pmQ7* strains (14 ± 0.8 days). The *coq7Δ/pDED* strain showed a slightly shorter mean CLS (11.4 ± 0.8 days) while the *coq7Δ/pAAA* strain had a clearly shorter mean CLS (9.1 ± 0.7 days). These results compared to CoQ₆ content (Figure 1A) support a model where the phosphorylation state of Coq7 affects CLS in yeast. Comparing the levels of CoQ₆ and CLS in the Coq7 versions we found an assortment of results. Thus, CoQ₆ level alone is not a factor that can explain the changes observed in CLS. Ptc7 regulates CoQ₆ biosynthesis by dephosphorylation of Coq7 [28]. We studied whether the lack of Ptc7 would re-




FIGURE 3: Chronological life span (CLS) is modified according to the phosphorylation state of Coq7. SDc-ura liquid cultures with 2% glucose as carbon source were inoculated at $0.1 \text{ OD}_{660\text{nm}}/\text{ml}$ with the indicated strains. After 5 days of growth, samples of each culture were harvested to measure the cell viability in YPD plates. The third day was used as control (100%). Plot shows a representative experiment repeated three times with similar results. See full dataset in Supplementary Material. **(A)** CLS in *COQ7* strains. Average life was *coq7Δ/pNMQ7* (12.1 ± 0.8), *coq7Δ/pRS316* (2.8 ± 0.2), *coq7Δ/pAAA* (9.1 ± 0.7), *coq7Δ/pDED* (11.4 ± 0.8) and *coq7Δ/pmQ7* (14 ± 0.8). **(B)** CLS in *PTC7* strains. Average life was BY4741 (12.7 ± 0.7), *ptc7Δ* (6.8 ± 0.4), *ptc7Δ + 10 μM CoQ₆* (6.9 ± 0.8) and *ptc7Δ:PTC7* (12.7 ± 0.5). **(C)** CoQ₆ of yeast strains analyzed in CLS experiment of *PTC7* gene strains. Similar data for *COQ7* gene CLS experiment was showed in Figure 1A. Cells from the indicated strains were subjected to mitochondrial purification, lipid extraction and quinone quantification. Samples were injected three times and the experiment was repeated at least three times.

sult in a compromised CLS. The *ptc7Δ* strain displayed a shortened mean CLS compared to wild type (6.8 ± 0.4 days versus 12.7 ± 0.7 days) (Figure 3B and Table S2). These results indicated that this phosphatase might regulate the normal function of longevity-associated pathways in yeast. CoQ₆ content of the *ptc7Δ* strain was significantly lower than both wild type and *ptc7Δ* strains rescued with the wild type allele (Figure 3C). As a result, CoQ₆ supplementation in CoQ₆ deficient *coq7Δ* yeast, rescues respiratory growth and oxidative stress resistance [45]. This was demonstrated by measuring the CLS of our strains in CoQ₆ supplemented media (Figure 3B). Interestingly, the addition of exogenous CoQ₆ to the *ptc7Δ* strain increased mitochondrial CoQ₆ to wild type levels (Figure 3C) but did not rescue CLS of the strain (7 ± 0.8 days), suggesting that the decreased CoQ₆ content is not responsible for the shortened CLS of the *ptc7Δ* strain. These results indicate that CoQ₆ levels cannot explain alterations of CLS in both *coq7Δ*

and *ptc7Δ* mutants. Moreover, the reduction in CLS in the *ptc7Δ* strain indicates that there might be additional functions of Ptc7 besides Coq7 activation.

Respiratory supercomplexes are altered in Coq7 phosphomutants

Mutant versions of Coq7 showed a clear defect on MRC activities focused in coupled reactions (NADH-cytochrome c reductase and succinate-cytochrome c reductase) that cannot be explained by the levels of CoQ₆. In the *coq7Δ/pAAA* strain the NADH-cytochrome c reductase activity is 40% less than wild type while the simple activity of both separated complexes is even higher than wild type. However, *coq7Δ/pAAA* mitochondria contain 250% of CoQ₆ compared to wild type (Figure 1). One possibility is that the expression of Coq7 versions can modify the stability or assembly of respiratory complexes. To this end, we analyzed the assembly of respiratory complexes by BN-PAGE in





FIGURE 4: Mitochondrial supercomplexes stability is affected in yeast expressing Coq7 phosphomutant versions. Pure mitochondrial samples obtained from the indicated strains were subjected to digitonin solubilization and 200 µg of each sample were analyzed by 1D BN-PAGE. (A) Coomassie staining of 1D BN-PAGE gels. Molecular weight markers are indicated on the left. On the right side, the three arrows indicate the calculated molecular weight of bands analyzed by densitometry in Figure 4C. (B) Upper panel, Western blot analysis blotted with antibody against the subunit Cox2 of Complex IV of a 1D BN-PAGE gel obtained in parallel to the Coomassie stained gel shown in Figure 4A using the same original mitochondrial digitonin solubilized samples. Lower panel, analysis by SDS-PAGE and Western blot of mitochondrial samples blotted with antibodies against the subunit Cox2 of Complex IV and porin. (C) Densitometry analysis of bands indicated with arrows in 4A. The analysis was performed with the software Image Lab 4.0 of Biorad. Band optical density was normalized with the full lane density and compared with the optical density of bands corresponding to the positive control, *coq7Δ/pNMQ7*. Results are representative of a set of two.

permeabilized mitochondria isolated from *coq7Δ* yeast expressing several versions of Coq7 (Figure 4). Cells were cultured first in YPD to increase cell mass and then the culture was transferred to YPG to activate mitochondrial metabolism. The *coq7Δ/pNMQ7* strain showed a typical profile of mitochondrial supercomplexes that mostly agrees with previously published data in wild type yeast [46], with three prominent bands at 911, 794 and 705 kDa, a double band around 480 kDa and a band located at 242 kDa (Figure 4A). The mass-spectrometry analysis of the three larger bands of *coq7Δ/pNMQ7* yeast indicated that the 911 kDa band corresponds mainly to complex V, the band of 794 kDa to complex III + complex IV and 705 KDa correspond to complex V. The detailed list of detected and identified proteins can be consulted at the Table S3 of Supplementary Material. Full MASCOT analysis is included in Supplementary Material. Immunoblot of this BN-PAGE gel with anti-Cox2 antibody (Figure 4B) showed differential intensities of supercomplex at 794 kDa but not in other bands. No reaction was observed in mitochondria from *coq7Δ/pDED* yeast, and very low in mitochondria from *coq7Δ/prS316* yeast. Lower intensities were also found in both *coq7Δ/pAAA* and *ptc7Δ* strains. Higher intensity was observed in control *coq7Δ/pNMQ7* and *coq7Δ/pmQ7* strains. The quantitative analysis of Coomassie staining showed that the 911 kDa and 794 kDa bands were significantly affected in both Coq7 mutants (Figure 4C). The expression of modified versions of Coq7 induced alterations in the assembly profile of respiratory complexes, being

more dramatic in the *coq7Δ/pDED* strain.

Mitophagy but not autophagy is affected in *ptc7Δ* mutant strains

Mitophagy is a general process that degrades damaged or non-useful mitochondria, which is required to maintain CLS in eukaryotic cells [31, 47, 48]. The *ptc7Δ* strain combines a low amount of CoQ₆ with a low CLS while the *coq7Δ/pDED* strain shows a low amount of CoQ₆ but a CLS comparable to wild type (Figures 1A and 3A). Both strains are equivalent in terms of Coq7 phosphorylation state and therefore Ptc7 must have another function independent of CoQ₆ biosynthesis that is responsible for the low CLS measured. We have previously demonstrated that the survival of human fibroblasts deficient in CoQ₁₀ production depends on a proper recycling of dysfunctional mitochondria by mitophagy [49]. We speculated that Ptc7 might regulate this process. Therefore, we analyzed whether the general process of autophagy and/or mitophagy could be compromised in *ptc7Δ*. Macroautophagy was analyzed by monitoring the Atg8p proteolysis using a plasmid expressing Atg8-GFP tag at the C-terminal under endogenous promoter (ATG8-GFP) (Figure 5A) [50]. Yeast cells were grown in YPD for 16 hours and subsequently were resuspended in nitrogen deprived medium (SDc-N) to induce macroautophagy. The positive control strain (BY4741; wild type) resulted in a marked increase of Atg8-GFP degradation, visible as free-GFP starting at 2 hours, which was maintained until 72 hours. According to previous reports, *pep4Δ* and *atg5Δ* strains,



GFP gene were grown in YPD for 16 hours. Then, yeasts were washed and media was replaced by YP Lactate. Protein extracts were analyzed using anti-GFP antibody for GFP-Om45, an outer membrane mitochondrial protein, and anti-Kar2 antibody for Kar2, an endoplasmic reticulum marker used as control for non-mitochondrial protein degradation. Original film plates used to prepare the panels are shown in Figure S1. Analysis of densitometry of panels B and C are shown in the Figure S2, S3 and S4. Data are mean ± SD, N ≥ 3 independent assays in all the experiments.

FIGURE 5: Mitophagy induction but no autophagy is compromised in Ptc7 yeast. (A) Yeast strains harboring an expression plasmid expressing the ATG8-GFP gene were grown in YPD 16 hours. Then, yeasts were washed and media was replaced by SDc-N with 2% glucose to activate autophagy. 0.5 OD_{660 nm} of Yeast cells were harvested at indicated time points. Protein extracts were analyzed by SDS-PAGE and Western blot using anti-GFP antibody. (B) Mitophagy analysis by porin degradation. Yeast strains were grown in YPD for 16 hours. Then, yeasts were washed and media was replaced by SDc-N with 2% glucose to activate mitophagy. Protein extracts were analyzed using anti-porin antibody. (C) Ptc7 overexpression increases the mitophagy induction. Yeast strains expressing the OM45-

which are deficient in protein degradation in the vacuole or autophagy respectively, showed impaired autophagy induction [51] (Figure 5A). In accordance with induced autophagy, a decrease of intact Atg8-GFP and a subsequent increase on free GFP was also observed in *ptc7Δ* strain, indicating that macroautophagy induction is not compromised in this strain.

To further study the potential role of Ptc7 in mitochondrial recycling, mitophagy was analyzed by monitoring the degradation of porin. Yeasts were grown in YPL for 16 hours and media was replaced with a nitrogen deprived media (SDc-N) to induce mitophagy (Figure 5B). The wild type strain showed a decreased amount of porin starting at day 2 and was gradually reduced until day 5. Porin degradation was not observed in the *ptc7Δ* strain under similar conditions, suggesting that Ptc7 is involved in mitochondrial recycling. Porin degradation was not observed in the autophagy-deficient yeast strains *atg5Δ* and *pep4Δ*. A protein expression analysis normalized with total protein loaded corroborated previous data (Figure 2S). To further determine the involvement of Ptc7 in mitophagy induction we determined the effect of Ptc7 overexpression on mitophagy (Figure 5C). Wild type yeast harboring the GFP protein fused in frame with the mitochondrial outer membrane protein Om45 were transformed with the empty yeast expression plasmid (pCM189) or containing the yeast *PTC7* coding sequence (pCM189-*PTC7*). Yeast cells were cultured in YPD for 16 hours and media was replaced with YPL, a non-fermentable carbon source, to induce mitochondrial biosynthesis. Yeast cells were grown through prolonged stationary phase to induce the selective recycling of mitochondria by mitophagy [31]. *PTC7* overexpression produced increased GFP free levels starting at 60 hours (270%) and at 120 hours (470%) of growth, indicating that the over-expression of *PTC7* enhances mitophagy induction (Figure S3). Remarkably, Kar2, a marker of endoplasmic reticulum, was not affected (Figure S4), indicating that macroautophagy was not activated by *PTC7* overexpression. Taken together, these data indicate that Ptc7 regulates or participates specifically in mitophagy but not in general macroautophagy, suggesting that this specific process may compromise CLS in *ptc7Δ* yeast.

DISCUSSION

Our understanding of the role of mitochondria on cell physiology and metabolism has evolved in the past decades from only a bioenergetics role to an interconnected organelle whose functions exceed energy supply. A key factor for part of these mitochondrial functions is CoQ, as proper regulation of its biosynthesis pathway exceeds cell bioenergetics and is tightly linked to cellular homeostasis. This idea supports the pleiotropic effect observed in patients with primary CoQ deficiency [7, 52]. Although most CoQ proteins are involved in enzymatic steps required for CoQ₆ biosynthesis [8], some CoQ proteins play a structural function and are required to stabilize the biosynthetic complex [8, 53]. In yeast, the shift from fermentation to respiration activates the expression of *COQ* genes to accommodate CoQ₆ biosynthesis to respiratory metabolism

[19, 54], which is associated to regulatory mechanisms such as biosynthesis complex assembly and post-translational modifications [9, 22, 25, 55].

The protein encoded by the *COQ7* gene has its catalytic activity controlled by complex regulatory mechanisms [16, 22, 25, 28, 56, 57]. Coq7 is a di-iron carboxylate protein with hydroxylase activity [58] that converts DMQ₆ in demethyl ubiquinone. The lack of Coq7 in yeast fully abolishes CoQ₆ biosynthesis and other intermediates although the expression of some point mutants such as e2519 (E223K) accumulates DMQ₆ [34], which would indicate that this step is a regulatory step in this pathway. DMQ₆ is also accumulated in wild type cells after the post-diauxic shift supporting this hypothesis [19, 34]. A model of CoQ₆ biosynthesis complex assembly, based on BN-PAGE, size exclusion chromatography and immunoprecipitation data, show that a precomplex of about 700 kDa is formed, which accumulates DMQ₆ [10] that, ultimately, will be converted into CoQ₆ after the recruitment of Coq7 to the complex [9, 55]. Coq7 is a phosphoprotein that, in the dephosphorylated state, activates CoQ₆ biosynthesis probably by interacting with the 700 kDa precomplex [22]. Here we have shown that the phosphomimetic Coq7 (Coq7-DED), which mimics Coq7 phosphorylation status in *ptc7Δ* strain [28], and the fully dephosphorylated Coq7 (Coq7-AAA) induced a decreased and an increased levels of CoQ₆ respectively, indicating that the regulatory step on CoQ₆ biosynthesis is at least partially controlled by Coq7 phosphorylation.

The physiological analysis of these strains confirms the catalytic function of Coq7 in CoQ₆ biosynthesis and its regulatory function in mitochondrial metabolism. Expression of both Coq7-AAA and Coq7-DED decreases antioxidant protection and increases the production of endogenous oxidative stress. A similar result was reported for the *ptc7Δ* strain [28]. These negative effects cannot be explained by the alteration of CoQ₆ levels, as the *coq7Δ/pmq7* strain, which also shows high levels of CoQ₆, had an endogenous oxidative stress and sensitivity similar to wild type. CoQ₆ has been reported as an antioxidant molecule mainly to protect cell membranes [35, 36, 59] but also as a pro-oxidant agent under physio-pathological conditions [60]. CoQ is involved in ROS production in the MRC mainly because of the transfer of electrons to complex III [61, 62]. It has been recently demonstrated in *Drosophila* that the reduced stage of CoQ (CoQH₂) causes CoQH₂-mediated ROS in complex I by retrograde electron transport and contributes to extend lifespan [63]. Similar conditions of higher CoQH₂ in mammal-cultured cells contribute to the partial degradation of complex I by the same mechanism [64]. Thus, we propose that the production of CoQ₆ in strains with low Coq7 phosphorylation are generating unbalanced CoQ₆ levels and redox stages out of the MRC complexes, which does not occur in the *coq7Δ/pmq7* strain. The unbalanced levels of CoQ₆ in these strains increase their sensitivity to external oxidative stress, decreasing their longevity. It has been reported previously that linolenic acid can induce mitochondrial oxidative stress [65]. The effect of linolenic acid may be enhanced by the mitochondrial dysfunction generated by the expression

of pAAA and pDED versions. In fact, the expression of pAAA or pDED induces a higher level of endogenous oxidative stress and also affects the stability of respiratory complexes, which possibly make those strains more susceptible to linolenic acid-induced oxidative stress.

CoQ is a component of the respirasome and it is proposed that pools of CoQ are bound to specific super assembly stages of respiratory complexes [66], which depends on the carbon source [64]. Here our data show that super-complexes are dissociated in yeast strains with unbalanced CoQ₆ concentrations, which also showed mitochondrial dysfunction, but not in the *COQ7* multicopy transformed yeast strain (*coq7Δ/pmQ7*) that exhibit a phenotype similar to wild type in both respiratory activities and assembly profile. Previous analyses of CoQ₆ biosynthesis complex showed that Coq7 is partially located in a large size complex but it can also be detected in smaller ones even as a monomer [10]. Coq7 interacts with Coq9, which shows lipid-binding activity and Coq7 interacting domains [16]. Several steps are required to integrate the CoQ₆ biosynthesis complex in yeast. First, there is an initial nucleation of Coq proteins around Coq4 [10, 67] to build up the Q-synthome, which requires stabilization by Coq8 [68], an unusual protein kinase that makes the pre-complex for-

mation [20] and demethoxy-Q₆ accumulation possible [19]. At this step Coq7 is mostly phosphorylated and it is not a component of the pre-complex [10]; Coq7 must then be dephosphorylated by Ptc7 to get activated and to bind to the fully active Q-synthome [9, 28, 55]. We speculate that CoQ₆ biosynthesis must be balanced with the components of the respiratory complexes. In light of our data, we hypothesize that the interaction of Coq7 with other proteins of the biosynthesis complex is a requirement for its integration in the MRC, and that extreme phosphorylation stages might prevent this possibility. This is supported by the data obtained from the *ptc7Δ* strain that mimics Coq7-DED expressing cells without eliminating all regulation of Coq7.

In fact, yeast contains two other mitochondrial phosphatases that belong to the same family of Ptc7, Ptc5 that participates in pyruvate dehydrogenase complex (PDH) regulation [30, 69] and Ptc6/Aup1 that participates in PDH regulation and is required for mitophagy activation [31, 70]. It is possible that these mitochondrial phosphatases may dephosphorylate Coq7 in the absence of Ptc7, which would explain the less severe effect on mitochondrial functions. Interestingly, the *ptc7Δ* strain shows shorten CLS compared to the *coq7Δ/pDED* strain and it is not rescued by CoQ₆ supplementation, although this strain shows a significant decrease of CoQ₆ content. CLS is only rescued when it is transformed with the homologous gene, which is also able to rescue the CoQ₆ content; overall these data indicate that Ptc7 has at least a dual function in yeast. As indicated above, Ptc6 is required for mitophagy activation [31, 70] and we have shown here that Ptc7 induces mitophagy as a mechanism to recycle defective mitochondria caused by CoQ₆ deficient MRC.

The lack of mitophagy in *ptc7Δ* yeast, which is a requisite to extend CLS [71, 72], can explain its shorten CLS and the negative recovery after CoQ₆ supplementation. We propose a dual and temporary differential function of Ptc7 on mitochondrial physiology and homeostasis (Figure 6). The entry of yeast on the post-diauxic shift (PDS) increases CoQ₆ biosynthesis and therefore respiratory metabolism by Coq7 activation. However, cell homeostasis during PDS requires the recycling of the excess and defective mitochondria. Ptc7, which would trigger mitophagy by dephosphorylating an unknown target involved in this process, would be a key regulator of mitochondria homeostasis in yeast by coordinating mitochondria recycling with CoQ₆ biosynthesis.




FIGURE 6: Model of Ptc7/Coq7 action to promote CLS extension. Ptc7 can regulate mitochondrial metabolisms in two different ways that are not directly related. After post diauxic shift, Coq7 expression and its activation by Ptc7 dephosphorylation leads to high levels of CoQ₆ concomitant with an increase of mitochondrial biogenesis. CLS extension requires a respiratory growth phase previous to the entrance in the stationary phase. The excess of mitochondria after the start of the stationary phase can be recycled by mitophagy to maintain an appropriated mitochondrial homeostasis. Ptc7 participates on mitophagy most likely by dephosphorylating an unknown target(s). Both functions, at different time frames, can help to promote a CLS extension in yeast.

MATERIAL AND METHODS

Yeast strains and growth media

Yeast strains used in this study are listed in Table 1. Growth media for yeast and bacteria were prepared as described previously [22]. Yeast cells were grown at 30°C with shaking (200 rpm).

Mitochondrial purification and BN-PAGE

Yeast cultures were grown in the appropriate culture media and mitochondria were purified according to the described method [73]. To solubilize mitochondria, 2 mg of pure mito-

TABLE 1. Strains used in this study.

Strains	Genotype	Source
coq7Δ	BY4741; MATα; his3Δ1; leu2Δ0; met15Δ0; ura3Δ0; YOR125c::KanMX4	Euroscarf
JM8	MAT α; Δade1; ρ^0	[74]
JM6	MAT a; Δhis4; ρ^0	[74]
ptc7Δ	BY4742; MATα; his3Δ1; leu2Δ0; lys2Δ0; ura3Δ0; YHR076w::KanMX4	Euroscarf
BY4742	MATα; his3Δ1; leu2Δ0; lys2Δ0; ura3Δ0.	Euroscarf
atg5Δ	BY4742, MATα; his3Δ1; leu2Δ0; lys2Δ0; ura3Δ0; YPL149w::kanMX4	Euroscarf
pep4Δ	BY4742; MATα; his3Δ1; leu2Δ0; lys2Δ0; ura3Δ0, YPL154c::kanMX4	Euroscarf
OM45::GFP	BY4741; MATα; his3Δ1; leu2Δ0; met15Δ0; ura3Δ0; YIL136W::GFP(S65T)-HIS3MX6	ATCC

chondria was incubated in 240 µl of solubilization buffer containing digitonin in a ratio 4:1 with protein, 1 mM PMSF, 10% glycerol, 150 mM potassium acetate and 30 mM HEPES, pH 7.4 for 30 min at 4°C. Solubilized samples were subjected to two rounds of centrifugation in a Beckman Coulter Microfuge 22R (15,000 × g, 15 min, at 4°C) and the supernatant was collected to BN-PAGE. Proteins quantification was performed by Bradford method (Biorad). BN-PAGE was performed with precast 3–12% gradient gels (NativePAGE™ Novex® Bis-Tris Gels) using the Xcell Sure Lock™ Mini-Cell electrophoresis system, including NativePAGE™ anode and cathode buffers according to the company instructions at 4°C. Lanes were loaded with 200 µg of mitochondrial solubilized supernatant and as MW marker NativeMark™ Unstained Protein Ladder was used. Gels were stained with Coomassie solution or were blotted onto PVDF, blocked in 5% Blocking Reagent (Biorad)

and phosphate-buffered saline. Proteins were detected by ECL using Luminata Crescendo (Millipore) and luminescence detected by Gel Doc XR+ image processing software (Bio-Rad). Cox2p (Novex Life Technologies), porin (Invitrogen) and Coq7 (Gift of Dr. C. F. Clarke, UCLA, USA) primary antibodies were used at 1: 2,000, 1: 1,000 and 1: 2,000 respectively. Goat anti-mouse and anti-rabbit secondary antibodies conjugated to horseradish peroxidase (Calbiochem) were used at a 1:5,000 dilutions.

Mitochondrial respiratory chain (MRC) activities

Fresh mitochondria were used to measure NADH-cytochrome c reductase, succinate-cytochrome c reductase activities and superoxide generation. Other MRC activities (NADH-DCIP reductase, succinate-DCIP reductase and decylubiquinol-cytochrome c reductase), were performed with fresh samples

TABLE 2. Vectors used in this study.

Name	Description	Specification	Source
pRS316	Yeast expression centromeric vector with <i>URA3</i> auxotrophy	Empty control	[34]
pRS426	Yeast expression episomic vector with <i>URA3</i> auxotrophy	Empty control	[34]
pNMQ7	pRS316 harboring the full yeast <i>COQ7</i> with promotor and terminator sequences	Positive control	[34]
pAAA	pRS316 with loss of function mutations in <i>COQ7</i> (S20A, S28A and S32A)	Non-phosphorylatable <i>COQ7</i> version	[22]
pDED	pRS316 with gain of function mutations in <i>COQ7</i> (S20D, S28E and S32D)	Mimicking a permanent phosphorylated <i>COQ7</i> version	[22]
pmQ7	pRS426 harboring yeast <i>COQ7</i>	<i>COQ7</i> multicopy complementation	[34]
ATG8-GFP	pRS316 harboring <i>ATG8</i> -GFP tag	<i>ATG8</i> tagged with GFP	[75]
pCM189	Yeast centromeric expression vector with <i>URA3</i> auxotrophy	Empty control	[76]
pCM189-PTC7	pCM189 harboring yeast <i>PTC7</i>	<i>PTC7</i> null mutant complementation	[28]

subjected to one freeze-thaw cycle. All MRC activities were determined according to previously published methods [34]. Superoxide generation was measured using the same method than for measuring complex III activity (decylubiquinol-cytochrome c reductase) but using acetylated cytochrome c instead cytochrome c [33]. H₂O₂ generation was performed using Amplex Red kit (Invitrogen) according to the manufacturer instructions.

Chronological Life Span (CLS)

Analysis was performed as previously described [37]. Briefly, cells were incubated in YPD and CLS was monitored starting at day 3 by quantification of colony forming units (CFUs) every 48 hours using the software OpenCFU 3.9 beta. The number of CFUs at day three was considered as 100% survival. Survival log-rank analyses Sigmastat 3.0 (SPSS) were calculated for each pair of lifespan analyses and average lifespan were shown in the corresponding dataset in Supplementary Material.

Protein mass spectrometry identification

Acrylamide gel bands were destained in NH₄HCO₃ 25 mM water/ACN 50:50. For reduction of cysteines, samples were incubated at 56°C for 60 min in 10 mM DTT (NH₄HCO₃ 25 mM). Cysteine carbamidomethylation was performed embedding bands in 55 mM IAA solution and incubating at room temperature for 30 min. Samples were in gel digested by trypsin (0.2 µg/µl in 1 mM HCl) diluted in NH₄HCO₃. Gel bands were covered with enzyme solution and incubated at 30°C overnight. Reaction was stopped with acetonitrile, and peptide were extracted adding 0.2% TFA. Prior to protein identification by MALDI-MS, we used nano-liquid chromatography for reversed phase peptide separation. Peptide fractioning was performed in a Proxeon EASY-nLC II apparatus with a C18 column (EASY-column, 75 µm x 100 mm) and mobile phases: Buffer A: 0.1% TFA (H₂O) and Buffer B: 0.1% TFA (ACN). Elution process was divided in the following flow steps: 0-48 min gradient 2%-45% B; 48-50 min gradient 45-100% B; 50-60 min isocratic 100% B. Fractions were collected every 15" using a Bruker Proteineer fc fraction collector and spotted onto a MALDI target plate. 192 samples were spotted and overlaid with 0.5 µl drops of HCCA matrix solution and left air dry. MALDI measurements were performed in a Bruker Ultraflexxtreme MALDI-TOF/TOF system, using Bruker Peptide Calibration standards as mass standards. A MALDI fingerprint spectrum was obtained for each fraction and peaks with higher intensity in each spot were selected as mass precursors for MS/MS peptide fragmentation experiments. Bruker WARP-LC software was used to process spectra and to integrate chromatography fractions data. Protein identification was carried out using MASCOT server. Ammonium bicarbonate, DL-dithiothreitol (DTT), iodoacetamide (IAA), trypsin from porcine pancreas, trifluoroacetic acid (TFA) and α-cyano-hydroxycinnamic acid (HCCA) were purchased from Sigma-Aldrich. Water (HPLC grade) and acetonitrile (HPLC grade) were purchased from Fluka. Peptide calibration standards were purchased from Bruker.

REFERENCES

- Doimo M, Desbats M a., Cerqua C, et al (2014). Genetics of Coenzyme Q10 Deficiency. *Mol Syndromol* 156–162.
- Emmanuele V, López LC, López L, et al (2012). Heterogeneity of coenzyme Q10 deficiency: patient study and literature review. *Arch Neurol* 69:978–83.

Total yeast protein extraction

Cells (10.10⁶ in 500 µl of water) were disrupted with 100 µl 2 M NaOH and 35% β-mercaptoethanol for 15 min on ice. Proteins were precipitated after the addition of 100 µl 3 M TCA for 15 min on ice. The pellet obtained after 15 min centrifugation on a microcentrifuge at full speed was washed with acetone, dried and resuspended in 30 µl of SDS-PAGE 1 x LB.

Other methods

CoQ₆ quantification was performed using mitochondrial samples according to previously published methods [28]. Densitometry analysis was carried out with a Gel Doc XR+ (Bio-Rad) with Image Lab 4.0 as software analysis. Statistical (t-Student) analyses were carried out using the Sigmastat 3.0 (SPSS) statistical package. Mitochondrial DNA integrity was checked in all strains using two ρ⁰ strains, JM6 and JM8 strains. All results are expressed as the average ± SD. Statistical analyses were carried out using the Sigmastat 3.0 (SPSS) statistical package.

ACKNOWLEDGMENTS

The research group is funded by the Andalusian Government as the BIO177 group through FEDER funds (European Commission), by the Ministerio de Economía y Competitividad, Instituto de Salud Carlos III (FIS PI14/01962) and by the International Q₁₀ Association.

AMMS received a predoctoral fellowship from the Consejería de Innovación Ciencia y Empresa, Junta de Andalucía (Spain). IGM received a predoctoral fellowship from the Plan Propio of the Universidad Pablo de Olavide de Sevilla.

The authors thank the group components for the critical reading of the manuscript and also to Ana Sanchez Cuesta for her technical help.

SUPPLEMENTAL MATERIAL

All supplemental data for this article are available online at www.microbialcell.com.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

COPYRIGHT

© 2017 González-Mariscal *et al.* This is an open-access article released under the terms of the Creative Commons Attribution (CC BY) license, which allows the unrestricted use, distribution, and reproduction in any medium, provided the original author and source are acknowledged.

Please cite this article as: I. González-Mariscal, A. Martín-Montalvo, C. Ojeda-González, A. Rodríguez-Eiguren, P. Gutiérrez-Ríos, P. Navas, and C. Santos-Ocaña (2017). Balanced CoQ₆ biosynthesis is required for lifespan and mitophagy in yeast. *Microbial Cell* 4(2): 38-51. doi: 10.15698/mic2017.02.556

3. DiMauro S, Quinzii CM, Hirano M (2007). Mutations in coenzyme Q10 biosynthetic genes. *J Clin Invest* 117:587–9.
4. Peng M, Falk MJ, Haase VH, et al (2008). Primary Coenzyme Q Deficiency in Pdss2 Mutant Mice Causes Isolated Renal Disease. *PLoS Genet* 4:e1000061.
5. Quinzii CM, Hirano M (2012). Primary and secondary CoQ(10) deficiencies in humans. *Biofactors* 37:361–5.
6. Yubero D, Montero R, Martín MA, et al (2016). Secondary coenzyme Q10 deficiencies in oxidative phosphorylation (OXPHOS) and non-OXPHOS disorders. *Mitochondrion* 30:51–58.
7. Santos-Ocaña C, Salvati L, Navas P (2015). The genes of CoQ10. In: Iain P. Hargreaves, Ph.D. (Neurometabolic Unit, National Hospital, Queen Square, London, UK) Assistant Editor: April K. Hargreaves, Ph.D. (Trinity University, Dublin I (ed) Coenzyme Q10 From fact to Fict., 1st ed. Nova Science Publisher, New York, pp 205–226
8. Tran UC, Clarke CF (2007). Endogenous synthesis of coenzyme Q in eukaryotes. *Mitochondrion* 7 Suppl:S62–71.
9. González-Mariscal I, García-Testón E, Padilla S, et al (2014). Regulation of coenzyme Q biosynthesis in yeast: A new complex in the block. *IUBMB Life* 66:63–70.
10. Marbois B, Gin P, Gulmezian M, Clarke CF (2009). The yeast Coq4 polypeptide organizes a mitochondrial protein complex essential for coenzyme Q biosynthesis. *Biochim Biophys Acta - Mol Cell Biol Lipids* 1791:69–75.
11. Leonard CJ, Aravind L, Koonin E V (1998). Novel families of putative protein kinases in bacteria and archaea: evolution of the "eukaryotic" protein kinase superfamily. *Genome Res* 8:1038–1047.
12. Macinga DR, Cook GM, Poole RK, Rather PN (1998). Identification and characterization of aarF, a locus required for production of ubiquinone in *Providencia stuartii* and *Escherichia coli* and for expression of 2'-N-acetyltransferase in *P. stuartii*. *J Bacteriol* 180:128–135.
13. Stefely JA, Reidenbach AG, Ulbrich A, et al (2015). Mitochondrial ADCK3 Employs an Atypical Protein Kinase-like Fold to Enable Coenzyme Q Biosynthesis. *Mol Cell* 57:83–94.
14. Stefely JA, Licitra F, Laredj L, et al (2016). Cerebellar Ataxia and Coenzyme Q Deficiency through Loss of Unorthodox Kinase Activity. *Mol Cell* 63:608–620.
15. Hsieh EJ, Gin P, Gulmezian M, et al (2007). *Saccharomyces cerevisiae* Coq9 polypeptide is a subunit of the mitochondrial coenzyme Q biosynthetic complex. *Arch Biochem Biophys* 463:19–26.
16. Lohman DC, Forouhar F, Beebe ET, et al (2014). Mitochondrial COQ9 is a lipid-binding protein that associates with COQ7 to enable coenzyme Q biosynthesis. *Proc Natl Acad Sci U S A*.
17. Gin P, Clarke CF (2005). Genetic evidence for a multi-subunit complex in coenzyme Q biosynthesis in yeast and the role of the Coq1 hexaprenyl diphosphate synthase. *J Biol Chem* 280:2676–81.
18. Marbois B, Gin P, Faull KF, et al (2005). Coq3 and Coq4 define a polypeptide complex in yeast mitochondria for the biosynthesis of coenzyme Q. *J Biol Chem* 280:20231–20238.
19. Padilla S, Tran UC, Jiménez-Hidalgo M, et al (2009). Hydroxylation of demethoxy-Q6 constitutes a control point in yeast coenzyme Q6 biosynthesis. *Cell Mol Life Sci* 66:173–86.
20. He CH, Xie LX, Allan CM, et al (2014). Coenzyme Q supplementation or over-expression of the yeast Coq8 putative kinase stabilizes multi-subunit Coq polypeptide complexes in yeast coq null mutants. *Biochim Biophys Acta* 1841:630–644.
21. Xie LX, Ozeir M, Tang JY, et al (2012). Over-expression of the Coq8 kinase in *Saccharomyces cerevisiae* coq null mutants allows for accumulation of diagnostic intermediates of the Coenzyme Q6 biosynthetic pathway. *J Biol Chem* 287:23571–81.
22. Martin-Montalvo A, Gonzalez-Mariscal I, Padilla S, et al (2011). Respiratory-induced coenzyme Q biosynthesis is regulated by a phosphorylation cycle of Cat5p/Coq7p. *Biochem J* 440:107–114.
23. Clarke CF (1996). The COQ7 Gene Encodes a Protein in *Saccharomyces cerevisiae* Necessary for Ubiquinone Biosynthesis. *J Biol Chem* 271:2995–3004.
24. Tauche A, Krause-Buchholz U, Rödel G, Rödel G (2008). Ubiquinone biosynthesis in *Saccharomyces cerevisiae*: the molecular organization of O-methylase Coq3p depends on Abc1p/Coq8p. *FEMS Yeast Res* 8:1263–75.
25. Xie LX, Hsieh EJ, Watanabe S, et al (2011). Expression of the human atypical kinase ADCK3 rescues coenzyme Q biosynthesis and phosphorylation of Coq polypeptides in yeast coq8 mutants. *Biochim Biophys Acta* 1811:348–360.
26. Busso C, Ferreira-Júnior JR, Paulela JA, et al (2015). Coq7p relevant residues for protein activity and stability. *Biochimie* 119:92–102.
27. Barford D, Das AK, Egloff MP (1998). The structure and mechanism of protein phosphatases: insights into catalysis and regulation. *Annu Rev Biophys Biomol Struct* 27:133–164.
28. Martin-Montalvo A, Gonzalez-Mariscal I, Pomares-Viciiana T, et al (2013). The Phosphatase Ptc7 Induces Coenzyme Q Biosynthesis by Activating the Hydroxylase Coq7 in Yeast. *J Biol Chem* 288:28126–28137.
29. Krause-Buchholz U, Gey U (2006). YIL042c and YOR090c encode the kinase and phosphatase of the *Saccharomyces cerevisiae* pyruvate dehydrogenase complex. *FEBS Lett* 580:2553–2560.
30. Gey U, Czupalla C, Hoflack B, et al (2008). Yeast pyruvate dehydrogenase complex is regulated by a concerted activity of two kinases and two phosphatases. *J Biol Chem* 283:9759–9767.
31. Tal R, Winter G, Ecker N, et al (2007). Aup1p, a yeast mitochondrial protein phosphatase homolog, is required for efficient stationary phase mitophagy and cell survival. *J Biol Chem* 282:5617–5624.
32. Ruan H, Yan Z, Sun H, Jiang L (2007). The YCR079w gene confers a rapamycin-resistant function and encodes the sixth type 2C protein phosphatase in *Saccharomyces cerevisiae*. *FEMS Yeast Res* 7:209–215.
33. Santos-Ocaña C, Villalba JM, Córdoba F, et al (1998). Genetic evidence for coenzyme Q requirement in plasma membrane electron transport. *J Bioenerg Biomembr* 30:465–75.
34. Padilla S, Jonassen T, Jiménez-Hidalgo MA, et al (2004). Demethoxy-Q, an intermediate of coenzyme Q biosynthesis, fails to support respiration in *Saccharomyces cerevisiae* and lacks antioxidant activity. *J Biol Chem* 279:25995–6004.
35. Poon WW, Do TQ, Noelle Marbois B, Clarke CF (1997). Sensitivity to treatment with polyunsaturated fatty acids is a general characteristic of the ubiquinone-deficient yeast coq mutants. *Mol Aspects Med* 18:121–127.
36. Do TQ, Schultz JR, Clarke CF (1996). Enhanced sensitivity of ubiquinone-deficient mutants of *Saccharomyces cerevisiae* to products of autoxidized polyunsaturated fatty acids. *Proc Natl Acad Sci U S A* 93:7534–9.
37. Fabrizio P, Longo VD (2003). The chronological life span of *Saccharomyces cerevisiae*. *Aging Cell* 2:73–81.

38. Barros MH, Bandy B, Tahara EB, Kowaltowski AJ (2004). Higher respiratory activity decreases mitochondrial reactive oxygen release and increases life span in *Saccharomyces cerevisiae*. *J Biol Chem* 279:49883–8.
39. Aerts AM, Zabrocki P, Govaert G, et al (2009). Mitochondrial dysfunction leads to reduced chronological lifespan and increased apoptosis in yeast. *FEBS Lett* 583:113–117.
40. Breitenbach M, Laun P, Dickinson JR, et al (2012). The role of mitochondria in the aging processes of yeast. *Subcell Biochem* 57:55–78.
41. Ocampo A, Liu J, Schroeder EA, et al (2012). Mitochondrial respiratory thresholds regulate yeast chronological life span and its extension by caloric restriction. *Cell Metab* 16:55–67.
42. Longo VD, Liou L-L, Valentine JS, Gralla EB (1999). Mitochondrial Superoxide Decreases Yeast Survival in Stationary Phase. *Arch Biochem Biophys* 365:131–142.
43. Fabrizio P, Liou L-L, Moy VN, et al (2003). SOD2 functions downstream of Sch9 to extend longevity in yeast. *Genetics* 163:35–46.
44. Gonidakis S, Longo VD (2008). Oxidative Stress and Aging in the Budding Yeast *Saccharomyces cerevisiae*. In: Miwa S, Beckman KB, Muller FL (eds) *Oxidative Stress Aging*. Humana Press, Totowa, NJ, pp 67–80.
45. Santos-Ocaña C, Do TQ, Padilla S, et al (2002). Uptake of exogenous coenzyme Q and transport to mitochondria is required for bc1 complex stability in yeast coq mutants. *J. Biol. Chem.* 277:
46. Schägger H, Pfeiffer K, Scha H, et al (2000). Supercomplexes in the respiratory chains of yeast and mammalian mitochondria. *EMBO J* 19:1777–83.
47. Youle RJ, Narendra DP (2011). Mechanisms of mitophagy. *Nat Rev Mol Cell Biol* 12:9–14.
48. Richard VR, Leonov A, Beach A, et al (2013). Macromitophagy is a longevity assurance process that in chronologically aging yeast limited in calorie supply sustains functional mitochondria and maintains cellular lipid homeostasis. *Aging (Albany NY)* 5:234–69.
49. Rodriguez-Hernandez A, Cordero MD, Salvati L, et al (2009). Coenzyme Q deficiency triggers mitochondria degradation by mitophagy. *Autophagy* 5:19–32.
50. Shintani T, Klionsky DJ (2004). Autophagy in health and disease: a double-edged sword. *Science* (80-) 306:990–995.
51. Kissova I, Deffieu M, Manon S, et al (2004). Uth1p is involved in the autophagic degradation of mitochondria. *J Biol Chem* 279:39068–39074.
52. Laredj LN, Licitra F, Puccio HM (2014). The molecular genetics of coenzyme Q biosynthesis in health and disease. *Biochimie* 100:78–87.
53. Kawamukai M (2016) Biosynthesis of coenzyme Q in eukaryotes. *Biosci Biotechnol Biochem* 80(1): 23–33.
54. Hagerman RA, Willis RA (2002). The yeast gene COQ5 is differentially regulated by Mig1p, Rtg3p and Hap2p. *Biochim Biophys Acta* 1578:51–58.
55. Allan CM, Awad AM, Johnson JS, et al (2015). Identification of Coq11, a new coenzyme Q biosynthetic protein in the CoQ-synthome in *Saccharomyces cerevisiae*. *J Biol Chem* jbc.M114.633131.
56. Brea-Calvo G, Siendones E, Sanchez-Alcazar JA, et al (2009). Cell survival from chemotherapy depends on NF-kappaB transcriptional up-regulation of coenzyme Q biosynthesis. *PLoS One* 4:e5301.
57. Cascajo M V, Abdelmohsen K, Noh JH, et al (2015). RNA-binding proteins regulate cell respiration and coenzyme Q biosynthesis by post-transcriptional regulation of COQ7. *RNA Biol* 0.
58. Stenmark P, Grünler J, Mattsson J, et al (2001). A new member of the family of di-iron carboxylate proteins. Coq7 (clk-1), a membrane-bound hydroxylase involved in ubiquinone biosynthesis. *J Biol Chem* 276:33297–300.
59. Maroz A, Anderson RF, Smith RA, Murphy MP (2009). Reactivity of ubiquinone and ubiquinol with superoxide and the hydroperoxyl radical: implications for in vivo antioxidant activity. *Free Radic Biol Med* 46:105–109.
60. Nohl H, Gille L, Schönheit K, Liu Y (1996). Conditions allowing redox-cycling ubisemiquinone in mitochondria to establish a direct redox couple with molecular oxygen. *Free Radic Biol Med* 20:207–213.
61. Boveris A, Cadena E, Stoppani AOCQ sintesi. y. funcio. (1976). Role of ubiquinone in the mitochondrial generation of hydrogen peroxide. *J Biol Chem* 156:435–444.
62. Turrens JF, Alexandre A, Lehninger AL (1985). Ubisemiquinone is the electron donor for superoxide formation by complex III of heart mitochondria. *Arch Biochem Biophys* 237:408–414.
63. Scialo F, Mallikarjun V, Stefanatos R, Sanz A (2013). Regulation of Lifespan by the Mitochondrial Electron Transport Chain: Reactive Oxygen Species-Dependent and Reactive Oxygen Species-Independent Mechanisms. *Antioxid Redox Signal* 19:1953–1969.
64. Guarás A, Perales-Clemente E, Calvo E, et al (2016). The CoQH₂/CoQ Ratio Serves as a Sensor of Respiratory Chain Efficiency. *Cell Rep* 15:197–209.
65. Colquhoun A, Schumacher RI (2001). gamma-Linolenic acid and eicosapentaenoic acid induce modifications in mitochondrial metabolism, reactive oxygen species generation, lipid peroxidation and apoptosis in Walker 256 rat carcinosarcoma cells. *Biochim Biophys Acta* 1533:207–219.
66. Lapuente-Brun E, Moreno-Loshuertos R, Acin-Perez R, et al (2013). Supercomplex Assembly Determines Electron Flux in the Mitochondrial Electron Transport Chain. *Science* (80-) 340:1567–1570.
67. González-Mariscal I, García-Testón E, Padilla S, et al (2014). The regulation of coenzyme q biosynthesis in eukaryotic cells: all that yeast can tell us. *Mol Syndromol* 5:107–18.
68. Do TQ, Hsu A Y, Jonassen T, et al (2001). A defect in coenzyme Q biosynthesis is responsible for the respiratory deficiency in *Saccharomyces cerevisiae* abc1 mutants. *J Biol Chem* 276:18161–8.
69. Ariño J, Casamayor A, González A, et al (2011). Type 2C protein phosphatases in fungi. *Eukaryot Cell* 10:21–33.
70. Journo D, Mor A, Abeliovich H (2009). Aup1-mediated regulation of Rtg3 during mitophagy. *J Biol Chem* 284:35885–35895.
71. Abeliovich H (2011). Stationary-Phase Mitophagy in Respiring *Saccharomyces cerevisiae*. *Antioxid Redox Signal* 14:2033–2011.
72. Sampaio-marques B, Burhans WC, Ludovico P (2014). Longevity pathways and maintenance of the proteome : the role of autophagy and mitophagy during yeast ageing. *Microb Cell* 1:118–127.
73. Glick BS, Pon LA (1995). Isolation of highly purified mitochondria from *Saccharomyces cerevisiae*. *Methods Enzymol* 260:213–223.
74. Proft M, Kötter P, Hedges D, et al (1995). CAT5, a new gene necessary for derepression of gluconeogenic enzymes in *Saccharomyces cerevisiae*. *EMBO J* 14:6116–26.
75. Shintani T, Klionsky DJ (2004). Cargo Proteins Facilitate the Formation of Transport Vesicles in the Cytoplasm to Vacuole Targeting Pathway. *J Biol Chem* 279:29889–29894.

76. Belli G, Gari E, Piedrafita L, et al (1998). An activator/repressor dual system allows tight tetracycline-regulated gene expression in budding yeast [published erratum appears in Nucleic Acids Res 1998 Apr 1;26(7):following 1855]. **Nucleic Acids Res** 26:942–947.